

Neutrino helicity flip by Cerenkov emission and absorption of plasmons in supernova

Subhendra Mohanty* and Sarira Sahu†

Physical Research Laboratory, Navrangpura, Ahmedabad - 380 009, India

We show that in a supernova core the longitudinal photon (plasmon) has a space-like dispersion and Cerenkov absorption and emission of such photons is kinematically allowed. If the neutrino has a non-zero magnetic moment, then helicity flipping Cerenkov absorption of a plasmon $\nu_L + \gamma \rightarrow \nu_R$ is the most efficient cooling mechanism of the supernova core, and this allows us to put a restrictive bound on the neutrino magnetic moment $\mu_\nu < 0.7 \times 10^{-13} \mu_B$.

It had been pointed out by Mohanty and Samal [1] that the helicity flipping Cerenkov process $\nu_L \rightarrow \nu_R + \gamma$ or $\gamma + \nu_L \rightarrow \nu_R$ could be an important cooling mechanism for the supernova core. Comparing the neutrino emissivity by the Cerenkov process with observations of SN1987A, a restrictive bound on the neutrino magnetic moment was established. Subsequently it was pointed out by Raffelt [2] that, this result was based on a numerical error in the calculation of the refractive index of the SN core and using the correct numbers it was shown that the photons in a SN core do not have a space-like dispersion relation, so the Cerenkov helicity flip process would not occur.

Here we show that the earlier estimate of refractive index was based on the thermodynamic formula for susceptibility which turns out to be invalid for real photons or plasmons even in the static limit. However an analysis of the dispersion relations of plasmons in an ultra relativistic plasma [3] shows that the longitudinal photon (plasmon) has a space-like branch and therefore in such a plasma the Cerenkov radiation of a plasmon is kinematically allowed [4]. We compute the neutrino emissivity of SN core by the Cerenkov helicity flip emission of a plasmon and subsequent escape of the ν_R . We show that the observations of neutrino flux from SN1987A put a constraint on the neutrino magnetic moment $\mu_\nu < 0.7 \times 10^{-13} \mu_B$.

The plasmon self energy diagram in a degenerate plasma is given by [3]

$$\Pi_L(k, w) = \frac{\tilde{\mu}_e^2 e^2}{\pi^2} \left(1 - \frac{w^2}{k^2} \right) \left[1 - \frac{w}{2k} \ln \left(\frac{w+k}{w-k} \right) \right] \quad (1)$$

where $\tilde{\mu}_e$ is the electron chemical potential which we have assumed to be larger than m_e and T . The dispersion relation for the longitudinal propagating mode $w^2 - k^2 = \text{Re } \Pi_L(k, w)$ is plotted in Figure 1. There is a lower branch which is space like. It is true that, there is also a large imaginary part of Π_L when $w^2 < k^2$, which means that this mode is Landau damped. However for our application all we need is that there must be an emission or an absorption of a plasmon accompanied by a neutrino

helicity flip.




Fig. 1. Dispersion relation for the degenerate plasma




Fig. 2. Plasmon refractive index as a function of frequency in a supernova core.

So it is not essential for our application that plasmon need to propagate far. Using this dispersion relation we find that in a supernova core there is a range of frequencies (0.3 – 50) MeV for which the refractive index $n = k/w > 1$ as shown in Figure 2 and therefore plasmon emission or absorption by the Cerenkov process is kinematically allowed in this range of plasmon frequencies.

The matrix element for the helicity flipping Cerenkov

process $\nu_L(p_1) \rightarrow \nu_R(p_2) + \gamma(k)$ is

$$\mathcal{M} = \mu_\nu \bar{u}(p_2) \sigma^{\mu\nu} u(p_1) \sqrt{Z_l} \epsilon_\mu(k, \lambda). \quad (2)$$

The photon polarisation sum in the medium for the longitudinal part is given by

$$\sum_\lambda \epsilon_\mu^L \epsilon_\nu^{L*} = \left(1 - \frac{1}{n^2}\right) u_\mu u_\nu + \frac{(u_\mu k_\nu + k_\mu u_\nu)}{n^2 w}, \quad (3)$$

where u_μ is the four velocity of the medium. The rate of energy emission by this process is

$$\dot{S} = \frac{\mu_\nu^2}{16\pi E_1^2} \int (E_1 - w) w^2 dw Z_l f(n) (w - 2E_1)^2 \quad (4)$$

where $f(n) = (n^2 - 1)^2/n^2$ and the wave function renormalisation factor Z_l is given by [3]

$$Z_l = \frac{2}{n^2 + \frac{3m_\gamma^2}{w^2} - 1}, \quad (5)$$

where $m_\gamma = e^2 \tilde{\mu}^2 / 3\pi^2$ is the effective photon mass in the medium. The neutrino luminosity is given by

$$Q_{\nu_R} = V \int_0^\infty 4\pi E_1^2 dE_1 f_\nu(E_1) \dot{S}(E_1), \quad (6)$$

where $f_\nu(E_1)$ is the distribution function of the neutrinos. Taking the SN core volume to be $V = 4.19 \times 10^{18} \text{ cm}^3$, the electron chemical potential $\tilde{\mu}_e = 280 \text{ MeV}$ and the neutrino chemical potential $\tilde{\mu}_\nu = 160 \text{ MeV}$ [5] we obtain the neutrino luminosity to be

$$Q_{\nu_R} = \mu_\nu^2 (1.45 \times 10^{61} \text{ MeV}^4), \quad (7)$$

in terms of the magnetic moment of the neutrino. Assuming that the entire energy of the core collapse is not carried away by the right handed neutrinos *i.e.*, $Q_{\nu_R} < 10^{52} \text{ ergs/sec}$ we obtain the upper bound on the neutrino magnetic moment $\mu_\nu < 0.2 \times 10^{-11} \mu_B$. This is comparable to the bound $(0.2 - 0.8) \times 10^{-11} \mu_B$ obtained [5] from the cooling of SN1987A by the helicity flip scattering.

At high temperature the rate of Cerenkov absorption process $\nu_L(p_1) + \gamma(k) \rightarrow \nu_R(p_2)$ is much larger than the Cerenkov emission. The cross section for the Cerenkov absorption is given by

$$\sigma(p_1, k) = \frac{\pi}{2E_1 w} \delta((p_1 + k)^2 - m_\nu^2) |\mathcal{M}|^2 \quad (8)$$

where

$$|\mathcal{M}|^2 = \mu^2 \frac{(n^2 - 1)^2}{n^2} Z_l w^2 (2E_1 + w)^2. \quad (9)$$

The rate of energy carried away by ν_R is given by

$$Q_{\nu_R} = V \int d^3 p_1 d^3 k f_\nu(p_1) f_\gamma(k) \sigma(p_1, k) (E_1 + w), \quad (10)$$

where $f_\nu(p_1)$ and $f_\gamma(k)$ are the distribution functions of the incoming ν_L and γ respectively. The expression for Q_{ν_R} can be written analytically as

$$Q_{\nu_R} = V 2\pi^3 \mu^2 T^7 \int_0^\infty dy \int_{x_1}^{x_2} dx g(x, y) Z_l \times \left(\frac{1}{e^{y - \tilde{\mu}_\nu/T} + 1} \right) \left(\frac{1}{e^x - 1} \right), \quad (11)$$

where the function $g(x, y)$ is defines as

$$g(x, y) = x^2 (x + y)(x + 2y)^2 (n^2 - 1)^2 \quad (12)$$

and $x = w/T$, $y = E_1/T$ are dimensionless quantities. The x integration is carried over the range where $n > 1$ and the Cerenkov absorption is kinematically allowed. Taking $\tilde{\mu}_{\nu_L} = 160 \text{ MeV}$, $T = 60 \text{ MeV}$ inside the supernova core [5] and doing the integral in Eq.(11) numerically we obtain the neutrino emissivity

$$Q_{\nu_R} = \mu_\nu^2 (1.1 \times 10^{64} \text{ MeV}^4) \quad (13)$$

in terms of the neutrino magnetic moment. Using the same constraint $Q_{\nu_R} < 10^{52} \text{ erg/sec}$ we obtain from Eq.(13) the upper bound on neutrino magnetic moment

$$\mu_\nu < 0.7 \times 10^{-13} \mu_B \quad (14)$$

This is two orders of magnitude lower than the earlier bounds [5]. This shows that if the neutrino has a non-zero magnetic moment then helicity flipping of the neutrinos by the Cerenkov absorption of plasmons is the most efficient cooling mechanism in a supernova.

Helicity flipping process in various types of plasmas have been studied using the quantum kinetic equation in ref [6].

* e-mail: mohanty@prl.ernet.in

† e-mail: sarira@prl.ernet.in

- [1] S. Mohanty and M. K. Samal, Phys. Rev. Lett. **77**, 806 (1996).
- [2] G. Raffelt, Phys. Rev. Lett. **79**, 773 (1997).
- [3] E. Bratten, Phys. Rev. Lett, **66**, 1655 (1991); H. A. Weldon, Phys. Rev. **D 26**, 1394 (1982); J. C. D'Olivo and J. F. Nieves, Nucleon effects on the photon diepersion relations in matter, hep-ph/9710305.
- [4] S. Sahu, Phys. Rev. **D 56**, 1688 (1997).
- [5] R. Barbieri and R. N. Mohapatra, Phys. Rev. Lett. **61**, 27 (1988).
- [6] Per Elmfors, Kari Enqvist, George Raffelt and Günter Sigl, Neutrinos with Magnetic moment: Depolarization rate in plasma, hep-ph/9703214.